

Medical Indications for Supplementation Word Search



Directions: Generate a list of words by answering the questions on the Medical Indications for Supplementation Question sheet. **Once you have completed your list, find and circle the words in the grid below.**

1. _____
2. _____
3. _____
4. _____
5. _____
6. _____

7. _____
8. _____
9. _____
10. _____
11. _____

L	G	O	M	O	T	H	E	R	S	T	R	I
T	N	E	M	E	V	O	M	L	E	W	O	B
E	I	N	F	E	C	T	I	O	N	S	S	M
R	D	R	O	E	E	A	V	M	E	A	S	S
E	E	E	S	D	I	N	L	K	I	I	L	O
E	E	G	E	B	I	O	S	L	L	E	H	E
A	F	A	B	T	A	I	O	O	I	E	N	A
L	T	I	O	T	R	T	B	E	E	I	N	E
C	S	N	O	I	T	A	C	I	D	E	M	K
O	A	S	I	N	T	T	E	O	A	M	E	C
H	E	R	P	E	S	S	I	M	P	L	E	X
O	R	M	M	N	T	E	M	A	E	A	I	I
L	B	F	T	I	T	G	E	E	O	A	E	D

Name: _____ Manager: _____

Please courier your completed sheet to **Susan Gallagher** at 1530 Markham Rd., 6th Floor or by fax to 416-338-6299.

Medical Indications for Supplementation Questions



*"Whenever interruption or cessation of breastfeeding is considered, the benefits of breastfeeding should be weighed against the risks posed by the use of human milk substitutes and the need to intervene because of the presenting medical condition."*¹

The following are conditions where the interruption, cessation or supplementation to breastfeeding may be required:

Infant Conditions:

- 1) Galactosemia, PKU, and Maple Syrup Urine Disease are inborn errors of _____ in infants that may require a partial or complete feeding with a breastmilk substitute.
- 2) Infants born weighing less than 1500g and born at less than 32 weeks _____ may need other food in addition to human milk for a limited period.
- 3) Temporary supplementation may be required for infants who are at _____ of hypoglycaemia because of impaired metabolic adaptation or increased glucose demand.
- 4) Infants with a significant weight loss and who fail to _____ birth weight by 2 weeks after birth may also require additional food for a short time.
- 5) Supplemental food may be needed in the interim for infants who demonstrate clinical indications of insufficient milk intake (such as no _____, or fewer than 1 a day in the first 2 weeks of life, or meconium 5 or more days after birth).

Maternal Conditions:

- 6) Sedating psychotherapeutic drugs, radioactive iodine, anti-epileptic drugs and opioids are _____ that are contraindicated to routine breastfeeding.
- 7) Cytotoxic chemotherapy requires that a mother stop _____ during therapy.
- 8) _____ should avoid excessive use of topical iodine, especially on open wounds or mucous membranes, when breastfeeding.
- 9) A temporary interruption of breastfeeding is recommended until all active _____ lesions found on the breast have been resolved.
- 10) Hepatitis B, hepatitis C, mastitis and breast abscesses are 4 maternal _____ that are not contraindicated for breastfeeding.
- 11) Even in situations of tobacco, _____, and drug use, breastfeeding remains the preferred feeding method for the majority of infants.

¹ Breastfeeding Committee for Canada. National Authority for the Baby-Friendly Initiative (BFI). BFI Integrated 10 Steps Practice Outcome Indicators for Hospitals and Community Health Services. www.breastfeedingcanada.ca/BFI